



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

deaths from cancer, 10 deaths from scarlet fever, 7 deaths from diphtheria, 6 deaths from measles, and 3 deaths from influenza. Finally, 4 persons died by violence.

The death rate of Berlin for the week ended June 10 was considerably higher than that for the preceding week and higher than it has been at any time during the past four months, amounting, calculated on the year, to 17.5 per thousand of the population, and was thus also higher than the rate for the same week of last year, in which it amounted to only 14.4 per thousand. In spite of the increased mortality of Berlin, however, at least one-half of the large German cities showed less favorable health conditions than this city, including Breslau, Königsberg, Rixdorf (with 28.9), Charlottenburg (with 20.4), Dresden, Leipzig, Halle, Magdeburg, Cologne, Frankfurt on the Main, Munich, and Stuttgart. On the other hand, the following cities had a lower death rate than Berlin, namely, Nuremberg, Düsseldorf, Hamburg, Bremen, Brunswick, Hanover, Schöneberg (with 11.1), and also London, while Paris and Vienna had a higher death rate than that of this city. The infant death rate rose from 4.5 per year and thousand in the preceding week to 5.7, being thereby somewhat higher than the Hamburg figure, but lower than the Munich and Leipzig rate. There were registered 3 cases of epidemic cerebro-spinal meningitis, 98 deaths from phthisis pulmonalis, 137 deaths from acute diseases of the respiratory organs, 39 deaths from cancer, 16 deaths from measles, 6 deaths from scarlet fever, 4 deaths from diphtheria, and 2 deaths from influenza. Finally, 3 persons died by violence.

GUATEMALA.

Report from Livingston, fruit port—Yellow fever.

Acting Assistant Surgeon Peters reports, June 24, as follows: Week ended June 24, 1905. Present officially estimated population, 3,500; 2 deaths from yellow fever; prevailing diseases, malarial and yellow fever; general sanitary condition of this port and the surrounding country during the week, fair.

Puerto Barrios so far remains free from infection.

Bills of health were issued to the following-named vessels:

Date.	Vessel.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of baggage disinfected.
June 21	Anselm	42
22	Anselm	42

The steamship Anselm took bills of health at both Puerto Barrios and Livingston.

HONDURAS.

Reports from Ceiba, fruit port.

Acting Assistant Surgeon Robertson reports as follows: Week ended June 24, 1905. Present officially estimated population, about 4,000; 1 death; prevailing diseases, malarial, mostly mild type; some severe forms; general sanitary condition of this port and the surrounding country during the week, good.